Imaging the source region of the 2003 San Simeon earthquake within the weak Franciscan Subduction Complex, Central California

Egill Hauksson, Caltech and David Oppenheimer and Tom Brocher, USGS


5th UJNR Earthquake Research Panel, October 12-16, 2004 Asilomar & Parkfield, California


Tasks


- •Determine 3-D *Vp* and *Vp/Vs* models
- •Refine hypocenters
- •Compare to finite source model of Chen Ji


Motivation


- •Relation to local fault structures
- Anomalous structure around the rupture
- Comparison with LA thrust earthquakes

























Methods and Approach


- 1) Minimum 1-D model


 VELEST from Kissling et al. (1993)
- 2) 3-D Vp and Vp/Vs Models
 Simulps from Thurber & Eberhart-Phillips (1993)
- 3) Relocations of hypocenters


 HypoDD Waldhauser & Ellsworth (2001)















Page et al., (1998)









Early Preliminary Relocations





Final Vp Model







Final Vp/Vs Model










Parkfield 2004



Major Late Quaternary Faults







Conclusions

- 1) Aftershocks reflect complex hanging wall deformation
- 2) The sequence occurred in the less rigid Franciscan formation
- 3) Similarities to Northridge and San Fernando but started at much shallower depth